

PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

OMIM

Books

Search PubMed



for



Limits

Preview/Index

History


Clipboard

Details

About Entrez



Abstract



Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: J Agric Food Chem 2002 Mar 27;50(7):2016-21

Related Articles, Books, LinkOut



Identification and quantification of aroma-active components that contribute to the distinct malty flavor of buckwheat honey.

Zhou Q, Wintersteen CL, Cadwallader KR.

Department of Food Science and Human Nutrition, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA.

Characteristic aroma components of buckwheat honey were studied by combined sensory and instrumental techniques. Relative aroma intensity of individual volatile components was evaluated by aroma extract dilution analysis (AEDA) of solvent extracts and by gas chromatography-olfactometry (GCO) of decreasing headspace samples (GCO-H). Results indicated that 3-methylbutanal, 3-hydroxy-4,5-dimethyl-2(5H)-furanone (sotolon), and (E)-beta-damascenone were the most potent odorants in buckwheat honey, with 3-methylbutanal being primarily responsible for the distinct malty aroma. Other important aroma-active compounds included methylpropanal, 2,3-butanedione, phenylacetaldehyde, 3-methylbutyric acid, maltol, vanillin, methional, coumarin, and p-cresol.

PMID: 11902950 [PubMed - indexed for MEDLINE]



Abstract



Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

sparc-sun-solaris2.8 May 24 2002 12:01:17